

Part 3: Trust-region methods for unconstrained optimization

Nick Gould (RAL)

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x)$$

MSc course on nonlinear optimization

LINESEARCH VS TRUST-REGION METHODS

- **Linesearch methods**
 - ◊ pick descent direction p_k
 - ◊ pick stepsize α_k to “reduce” $f(x_k + \alpha p_k)$
 - ◊ $x_{k+1} = x_k + \alpha_k p_k$
- **Trust-region methods**
 - ◊ pick step s_k to reduce “model” of $f(x_k + s)$
 - ◊ accept $x_{k+1} = x_k + s_k$ if decrease in model inherited by $f(x_k + s_k)$
 - ◊ otherwise set $x_{k+1} = x_k$, “refine” model

UNCONSTRAINED MINIMIZATION

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x)$$

where the **objective function** $f : \mathbb{R}^n \rightarrow \mathbb{R}$

- assume that $f \in C^1$ (sometimes C^2) and Lipschitz
- often in practice this assumption violated, but not necessary

TRUST-REGION MODEL PROBLEM

Model $f(x_k + s)$ by:

- linear model
$$m_k^L(s) = f_k + s^T g_k$$
- quadratic model — symmetric B_k
$$m_k^Q(s) = f_k + s^T g_k + \frac{1}{2} s^T B_k s$$

Major difficulties:

- models may not resemble $f(x_k + s)$ if s is large
- models may be unbounded from below
 - ◊ linear model - always unless $g_k = 0$
 - ◊ quadratic model - always if B_k is indefinite, possibly if B_k is only positive semi-definite

THE TRUST REGION

Prevent model $m_k(s)$ from unboundedness by imposing a **trust-region** constraint

$$\|s\| \leq \Delta_k$$

for some “suitable” scalar **radius** $\Delta_k > 0$

⇒ **trust-region subproblem**

approx minimize $m_k(s)$ subject to $\|s\| \leq \Delta_k$
 $s \in \mathbb{R}^n$

- in theory does not depend on norm $\|\cdot\|$
- in practice it might!

BASIC TRUST-REGION METHOD

Given $k = 0$, $\Delta_0 > 0$ and x_0 , until “convergence” do:

Build the second-order model $m(s)$ of $f(x_k + s)$.

“Solve” the trust-region subproblem to find s_k

for which $m(s_k)$ “<” f_k and $\|s_k\| \leq \Delta_k$, and define

$$\rho_k = \frac{f_k - f(x_k + s_k)}{f_k - m_k(s_k)}.$$

If $\rho_k \geq \eta_0$ [**very successful**]

set $x_{k+1} = x_k + s_k$ and $\Delta_{k+1} = \gamma_l \Delta_k$

Otherwise if $\rho_k \geq \eta_s$ then [**successful**]

set $x_{k+1} = x_k + s_k$ and $\Delta_{k+1} = \Delta_k$

Otherwise [**unsuccessful**]

set $x_{k+1} = x_k$ and $\Delta_{k+1} = \gamma_d \Delta_k$

Increase k by 1

$$\boxed{0 < \eta_0 < 1}$$

$$\boxed{\gamma_l \geq 1}$$

$$\boxed{0 < \eta_s \leq \eta_0 < 1}$$

$$\boxed{0 < \gamma_d < 1}$$

OUR MODEL

For simplicity, concentrate on the second-order (Newton-like) model

$$m_k(s) = m_k^Q(s) = f_k + s^T g_k + \frac{1}{2} s^T B_k s$$

and the ℓ_2 -trust region norm $\|\cdot\| = \|\cdot\|_2$

Note:

- $B_k = H_k$ is allowed
- analysis for other trust-region norms simply adds extra constants in following results

“SOLVE” THE TRUST REGION SUBPROBLEM?

At the very least

- aim to achieve as much reduction in the model as would an iteration of steepest descent

- **Cauchy point**: $s_k^C = -\alpha_k^C g_k$ where

$$\begin{aligned} \alpha_k^C &= \arg \min_{\alpha > 0} m_k(-\alpha g_k) \text{ subject to } \alpha \|g_k\| \leq \Delta_k \\ &= \arg \min_{0 < \alpha \leq \Delta_k / \|g_k\|} m_k(-\alpha g_k) \end{aligned}$$

◊ minimize quadratic on line segment ⇒ very easy!

- require that

$$m_k(s_k) \leq m_k(s_k^C) \text{ and } \|s_k\| \leq \Delta_k$$

- in practice, hope to do far better than this

ACHIEVABLE MODEL DECREASE

Theorem 3.1. If $m_k(s)$ is the second-order model and s_k^c is its Cauchy point within the trust-region $\|s\| \leq \Delta_k$,

$$f_k - m_k(s_k^c) \geq \frac{1}{2} \|g_k\| \min \left[\frac{\|g_k\|}{1 + \|B_k\|}, \Delta_k \right].$$

PROOF OF THEOREM 3.1

$$m_k(-\alpha g_k) = f_k - \alpha \|g_k\|^2 + \frac{1}{2} \alpha^2 g_k^T B_k g_k.$$

Result immediate if $g_k = 0$.

Otherwise, 3 possibilities

- (i) curvature $g_k^T B_k g_k \leq 0 \implies m_k(-\alpha g_k)$ unbounded from below as α increases \implies Cauchy point occurs on the trust-region boundary.
- (ii) curvature $g_k^T B_k g_k > 0 \&$ minimizer $m_k(-\alpha g_k)$ occurs at or beyond the trust-region boundary \implies Cauchy point occurs on the trust-region boundary.
- (iii) the curvature $g_k^T B_k g_k > 0 \&$ minimizer $m_k(-\alpha g_k)$, and hence Cauchy point, occurs before trust-region is reached.

Consider each case in turn;

Case (i)

$$g_k^T B_k g_k \leq 0 \& \alpha \geq 0 \implies$$

$$m_k(-\alpha g_k) = f_k - \alpha \|g_k\|^2 + \frac{1}{2} \alpha^2 g_k^T B_k g_k \leq f_k - \alpha \|g_k\|^2 \quad (1)$$

Cauchy point lies on boundary of the trust region \implies

$$\alpha_k^c = \frac{\Delta_k}{\|g_k\|}. \quad (2)$$

$$(1) + (2) \implies$$

$$f_k - m_k(s_k^c) \geq \|g_k\|^2 \frac{\Delta_k}{\|g_k\|} = \|g_k\| \Delta_k \geq \frac{1}{2} \|g_k\| \Delta_k.$$

Case (ii)

$$\alpha_k^* \stackrel{\text{def}}{=} \arg \min m_k(-\alpha g_k) \equiv f_k - \alpha \|g_k\|^2 + \frac{1}{2} \alpha^2 g_k^T B_k g_k \quad (3)$$

$$\implies$$

$$\alpha_k^* = \frac{\|g_k\|^2}{g_k^T B_k g_k} \geq \alpha_k^c = \frac{\Delta_k}{\|g_k\|} \quad (4)$$

$$\implies$$

$$\alpha_k^c g_k^T B_k g_k \leq \|g_k\|^2. \quad (5)$$

$$(3) + (4) + (5) \implies$$

$$\begin{aligned} f_k - m_k(s_k^c) &= \alpha_k^c \|g_k\|^2 - \frac{1}{2} [\alpha_k^c]^2 g_k^T B_k g_k \geq \frac{1}{2} \alpha_k^c \|g_k\|^2 \\ &= \frac{1}{2} \|g_k\|^2 \frac{\Delta_k}{\|g_k\|} = \frac{1}{2} \|g_k\| \Delta_k. \end{aligned}$$

Case (iii)

$$\alpha_k^c = \alpha_k^* = \frac{\|g_k\|^2}{g_k^T B_k g_k}$$

\implies

$$\begin{aligned} f_k - m_k(s_k^c) &= \alpha_k^* \|g_k\|^2 + \frac{1}{2} (\alpha_k^*)^2 g_k^T B_k g_k \\ &= \frac{\|g_k\|^4}{\|g_k\|^4} - \frac{1}{2} \frac{\|g_k\|^4}{\|g_k\|^4} \\ &= \frac{g_k^T B_k g_k}{\|g_k\|^4} - \frac{1}{2} \frac{g_k^T B_k g_k}{\|g_k\|^4} \\ &= \frac{1}{2} \frac{g_k^T B_k g_k}{\|g_k\|^4} \\ &\geq \frac{1}{2} \frac{1}{\|B_k\|}, \end{aligned}$$

where

$$|g_k^T B_k g_k| \leq \|g_k\|^2 \|B_k\| \leq \|g_k\|^2 (1 + \|B_k\|)$$

because of the Cauchy-Schwarz inequality.

DIFFERENCE BETWEEN MODEL AND FUNCTION

Lemma 3.3. Suppose that $f \in C^2$, and that the true and model Hessians satisfy the bounds $\|H(x)\| \leq \kappa_h$ for all x and $\|B_k\| \leq \kappa_b$ for all k and some $\kappa_h \geq 1$ and $\kappa_b \geq 0$. Then

$$|f(x_k + s_k) - m_k(s_k)| \leq \kappa_d \Delta_k^2,$$

where $\kappa_d = \frac{1}{2}(\kappa_h + \kappa_b)$, for all k .

Corollary 3.2. If $m_k(s)$ is the second-order model, and s_k is an improvement on the Cauchy point within the trust-region $\|s\| \leq \Delta_k$,

$$f_k - m_k(s_k) \geq \frac{1}{2} \|g_k\| \min \left[\frac{\|g_k\|}{1 + \|B_k\|}, \Delta_k \right].$$

PROOF OF LEMMA 3.3

Mean value theorem \implies

$$f(x_k + s_k) = f(x_k) + s_k^T \nabla_x f(x_k) + \frac{1}{2} s_k^T \nabla_{xx} f(\xi_k) s_k$$

for some $\xi_k \in [x_k, x_k + s_k]$. Thus

$$\begin{aligned} |f(x_k + s_k) - m_k(s_k)| &= \frac{1}{2} |s_k^T H(\xi_k) s_k - s_k^T B_k s_k| \leq \frac{1}{2} |s_k^T H(\xi_k) s_k| + \frac{1}{2} |s_k^T B_k s_k| \\ &\leq \frac{1}{2} (\kappa_h + \kappa_b) \|s_k\|^2 \leq \kappa_d \Delta_k^2 \end{aligned}$$

using the triangle and Cauchy-Schwarz inequalities.

ULTIMATE PROGRESS AT NON-OPTIMAL POINTS

Lemma 3.4. Suppose that $f \in C^2$, that the true and model Hessians satisfy the bounds $\|H_k\| \leq \kappa_h$ and $\|B_k\| \leq \kappa_b$ for all k and some $\kappa_h \geq 1$ and $\kappa_b \geq 0$, and that $\kappa_d = \frac{1}{2}(\kappa_h + \kappa_b)$. Suppose furthermore that $g_k \neq 0$ and that

$$\Delta_k \leq \|g_k\| \min \left(\frac{1}{\kappa_h + \kappa_b}, \frac{(1 - \eta_v)}{2\kappa_d} \right).$$

Then iteration k is very successful and

$$\Delta_{k+1} \geq \Delta_k.$$

PROOF OF LEMMA 3.4

By definition,

$$1 + \|B_k\| \leq \kappa_h + \kappa_b$$

+ first bound on $\Delta_k \implies$

$$\Delta_k \leq \frac{\|g_k\|}{\kappa_h + \kappa_b} \leq \frac{\|g_k\|}{1 + \|B_k\|}.$$

Corollary 3.2 \implies

$$f_k - m_k(s_k) \geq \frac{1}{2}\|g_k\| \min \left[\frac{\|g_k\|}{1 + \|B_k\|}, \Delta_k \right] = \frac{1}{2}\|g_k\| \Delta_k.$$

+ Lemma 3.3 + second bound on $\Delta_k \implies$

$$|p_k - 1| = \left| \frac{f(x_k + s_k) - m_k(s_k)}{f_k - m_k(s_k)} \right| \leq 2 \frac{\kappa_d \Delta_k^2}{\|g_k\| \Delta_k} = 2 \frac{\kappa_d \Delta_k}{\|g_k\|} \leq 1 - \eta_v.$$

$\implies p_k \geq \eta_v \implies$ iteration is very successful.

RADIUS WON'T SHRINK TO ZERO AT NON-OPTIMAL POINTS

Lemma 3.5. Suppose that $f \in C^2$, that the true and model Hessians satisfy the bounds $\|H_k\| \leq \kappa_h$ and $\|B_k\| \leq \kappa_b$ for all k and some $\kappa_h \geq 1$ and $\kappa_b \geq 0$, and that $\kappa_d = \frac{1}{2}(\kappa_h + \kappa_b)$. Suppose furthermore that there exists a constant $\epsilon > 0$ such that $\|g_k\| \geq \epsilon$ for all k . Then

$$\Delta_k \geq \kappa_\epsilon \stackrel{\text{def}}{=} \epsilon \gamma_d \min \left(\frac{1}{\kappa_h + \kappa_b}, \frac{(1 - \eta_v)}{2\kappa_d} \right)$$

for all k .

PROOF OF LEMMA 3.5

Suppose otherwise that iteration k is first for which

$$\Delta_{k+1} \leq \kappa_\epsilon.$$

$\Delta_k > \Delta_{k+1} \implies$ iteration k unsuccessful $\implies \gamma_d \Delta_k \leq \Delta_{k+1}$. Hence

$$\begin{aligned} \Delta_k &\leq \epsilon \min \left(\frac{1}{\kappa_h + \kappa_b}, \frac{(1 - \eta_v)}{2\kappa_d} \right) \\ &\leq \|g_k\| \min \left(\frac{1}{\kappa_h + \kappa_b}, \frac{(1 - \eta_v)}{2\kappa_d} \right) \end{aligned}$$

But this contradicts assertion of Lemma 3.4 that iteration k must be very successful.

POSSIBLE FINITE TERMINATION

Lemma 3.6. Suppose that $f \in C^2$, and that both the true and model Hessians remain bounded for all k . Suppose furthermore that there are only finitely many successful iterations. Then $x_k = x_*$ for all sufficiently large k and $g(x_*) = 0$.

PROOF OF LEMMA 3.6

$$x_{k_0+j} = x_{k_0+1} = x_*$$

for all $j > 0$, where k_0 is index of last successful iterate.

All iterations are unsuccessful for sufficiently large $k \implies \{\Delta_k\} \longrightarrow 0$ + Lemma 3.4 then implies that if $\|g_{k_0+1}\| > 0$ there must be a successful iteration of index larger than k_0 , which is impossible $\implies \|g_{k_0+1}\| = 0$.

GLOBAL CONVERGENCE OF ONE SEQUENCE

Theorem 3.7. Suppose that $f \in C^2$, and that both the true and model Hessians remain bounded for all k . Then either

$$g_l = 0 \text{ for some } l \geq 0$$

or

$$\lim_{k \rightarrow \infty} f_k = -\infty$$

or

$$\liminf_{k \rightarrow \infty} \|g_k\| = 0.$$

PROOF OF THEOREM 3.7

Let \mathbf{S} be the index set of successful iterations. Lemma 3.6 \implies true Theorem 3.7 when $|\mathbf{S}|$ finite.

So consider $|\mathbf{S}| = \infty$, and suppose f_k bounded below and

$$\|g_k\| \geq \epsilon \tag{6}$$

for some $\epsilon > 0$ and all k , and consider some $k \in \mathbf{S}$.

+ Corollary 3.2, Lemma 3.5, and the assumption (6) \implies

$$f_k - f_{k+1} \geq \eta_{\mathbf{S}} [f_k - m_k(s_k)] \geq \delta_\epsilon \stackrel{\text{def}}{=} \frac{1}{2} \eta_{\mathbf{S}} \epsilon \min \left[\frac{\epsilon}{1 + \kappa_b}, \kappa_\epsilon \right].$$

\implies

$$f_0 - f_{k+1} = \sum_{\substack{j=0 \\ j \in \mathbf{S}}}^k [f_j - f_{j+1}] \geq \sigma_k \delta_\epsilon,$$

where σ_k is the number of successful iterations up to iteration k . But

$$\lim_{k \rightarrow \infty} \sigma_k = +\infty.$$

$\implies f_k$ unbounded below \implies a subsequence of the $\|g_k\| \longrightarrow 0$

GLOBAL CONVERGENCE

Theorem 3.8. Suppose that $f \in C^2$, and that both the true and model Hessians remain bounded for all k . Then either

$$g_l = 0 \text{ for some } l \geq 0$$

or

$$\lim_{k \rightarrow \infty} f_k = -\infty$$

or

$$\lim_{k \rightarrow \infty} g_k = 0.$$

II: SOLVING THE TRUST-REGION SUBPROBLEM

(approximately) minimize $q(s) \equiv s^T g + \frac{1}{2} s^T B s$ subject to $\|s\| \leq \Delta$
 $s \in \mathbb{R}^n$

AIM: find s_* so that

$$q(s_*) \leq q(s^c) \text{ and } \|s_*\| \leq \Delta$$

Might solve

- exactly \implies Newton-like method
- approximately \implies steepest descent/conjugate gradients

THE ℓ_2 -NORM TRUST-REGION SUBPROBLEM

minimize $q(s) \equiv s^T g + \frac{1}{2} s^T B s$ subject to $\|s\|_2 \leq \Delta$
 $s \in \mathbb{R}^n$

Solution characterisation result:

Theorem 3.9. Any *global* minimizer s_* of $q(s)$ subject to $\|s\|_2 \leq \Delta$ satisfies the equation

$$(B + \lambda_* I) s_* = -g,$$

where $B + \lambda_* I$ is positive semi-definite, $\lambda_* \geq 0$ and $\lambda_* (\|s_*\|_2 - \Delta) = 0$. If $B + \lambda_* I$ is positive definite, s_* is unique.

PROOF OF THEOREM 3.9

Problem equivalent to minimizing $q(s)$ subject to $\frac{1}{2} \Delta^2 - \frac{1}{2} s^T s \geq 0$.
 Theorem 1.9 \implies

$$g + B s_* = -\lambda_* s_* \tag{7}$$

for some Lagrange multiplier $\lambda_* \geq 0$ for which either $\lambda_* = 0$ or $\|s_*\|_2 = \Delta$ (or both). It remains to show $B + \lambda_* I$ is positive semi-definite.

If s_* lies in the interior of the trust-region, $\lambda_* = 0$, and Theorem 1.10 $\implies B + \lambda_* I = B$ is positive semi-definite.

If $\|s_*\|_2 = \Delta$ and $\lambda_* = 0$, Theorem 1.10 $\implies v^T B v \geq 0$ for all $v \in \mathcal{N}_+ = \{v | s_*^T v \geq 0\}$. If $v \notin \mathcal{N}_+ \implies -v \in \mathcal{N}_+ \implies v^T B v \geq 0$ for all v .

Only remaining case is where $\|s_*\|_2 = \Delta$ and $\lambda_* > 0$. Theorem 1.10 $\implies v^T (B + \lambda_* I) v \geq 0$ for all $v \in \mathcal{N}_+ = \{v | s_*^T v = 0\} \implies$ remains to consider $v^T B v$ when $s_*^T v \neq 0$.




Figure 3.1: Construction of "missing" directions of positive curvature.

ALGORITHMS FOR THE ℓ_2 -NORM SUBPROBLEM

Two cases:

- B positive-semi definite and $Bs = -g$ satisfies $\|s\|_2 \leq \Delta \implies s_* = s$
- B indefinite or $Bs = -g$ satisfies $\|s\|_2 > \Delta$
In this case
 - ◊ $(B + \lambda_* I)s_* = -g$ and $s_*^T s_* = \Delta^2$
 - ◊ nonlinear (quadratic) system in s and λ
 - ◊ concentrate on this

Let s be any point on the boundary δR of the trust-region R , and let $w = s - s_*$. Then

$$-w^T s_* = (s_* - s)^T s_* = \frac{1}{2}(s_* - s)^T (s_* - s) = \frac{1}{2}w^T w \quad (8)$$

since $\|s\|_2 = \Delta = \|s_*\|_2$. (7) + (8) \implies

$$\begin{aligned} q(s) - q(s_*) &= w^T (g + Bs_*) + \frac{1}{2}w^T Bw \\ &= -\lambda_* w^T s_* + \frac{1}{2}w^T Bw = \frac{1}{2}w^T (B + \lambda_* I)w, \end{aligned} \quad (9)$$

$\implies w^T (B + \lambda_* I)w \geq 0$ since s_* is a global minimizer. But

$$s = s_* - 2 \frac{s_*^T v}{q^T v} v \in \delta R$$

\implies (for this s) $w\|v \implies v^T (B + \lambda_* I)v \geq 0$.

When $B + \lambda_* I$ is positive definite, $s_* = -(B + \lambda_* I)^{-1}g$. If $s_* \in \delta R$ and $s \in R$, (8) and (9) become $-w^T s_* \geq \frac{1}{2}w^T w$ and $q(s) \geq q(s_*) + \frac{1}{2}w^T (B + \lambda_* I)w$ respectively. Hence, $q(s) > q(s_*)$ for any $s \neq s_*$. If s_* is interior, $\lambda_* = 0$, B is positive definite, and thus s_* is the unique unconstrained minimizer of $q(s)$.

EQUALITY CONSTRAINED ℓ_2 -NORM SUBPROBLEM

Suppose B has spectral decomposition

$$B = U^T \Lambda U$$

- U eigenvectors
 - Λ diagonal eigenvalues: $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$
- Require $B + \lambda I$ positive semi-definite $\implies \lambda \geq -\lambda_1$

Define

$$s(\lambda) = -(B + \lambda I)^{-1}g$$


Require

$$\psi(\lambda) \stackrel{\text{def}}{=} \|s(\lambda)\|_2^2 = \Delta^2$$


Note

$$\begin{aligned} \psi(\lambda) &= \|U^T (\Lambda + \lambda I)^{-1} U g\|_2^2 = \sum_{i=1}^n \frac{\gamma_i^2}{(\lambda_i + \lambda)^2} \\ &\quad (\gamma_i = e_i^T U g) \end{aligned}$$


CONVEX EXAMPLE



NONCONVEX EXAMPLE



THE "HARD" CASE



SUMMARY

For indefinite B ,

Hard case occurs when g orthogonal to eigenvector u_1 for most negative eigenvalue λ_1

- OK if radius is radius small enough
- No "obvious" solution to equations ... but solution is actually of the form

$$s_{\text{lim}} + \sigma u_1$$

where

- ◊ $s_{\text{lim}} = \lim_{\lambda \rightarrow -\lambda_1} s(\lambda)$
- ◊ $\|s_{\text{lim}} + \sigma u_1\|_2 = \Delta$

HOW TO SOLVE $\|s(\lambda)\|_2 = \Delta$

DON'T!!


Solve instead the **secular equation**

$$\phi(\lambda) \stackrel{\text{def}}{=} \frac{1}{\|s(\lambda)\|_2} - \frac{1}{\Delta} = 0$$

- no poles
- smallest at eigenvalues (except in hard case!)
- analytic function \implies ideal for Newton
- global convergent (ultimately quadratic rate except in hard case)
- need to safeguard to protect Newton from the hard & interior solution cases

THE SECULAR EQUATION

$$\begin{aligned} &\min -\frac{1}{4}s_1^2 + \frac{1}{4}s_2^2 + \frac{1}{2}s_1 + s_2 \\ &\text{subject to } \|s\|_2 \leq 4 \end{aligned}$$



NEWTON'S METHOD FOR SECULAR EQUATION

Newton correction at λ is $-\phi(\lambda)/\phi'(\lambda)$. Differentiating

$$\begin{aligned} \phi(\lambda) &= \frac{1}{\|s(\lambda)\|_2} - \frac{1}{\Delta} = \frac{1}{(s^T(\lambda)s(\lambda))^{\frac{1}{2}}} - \frac{1}{\Delta} \implies \\ \phi'(\lambda) &= -\frac{s^T(\lambda)\nabla_{\lambda}s(\lambda)}{(s^T(\lambda)s(\lambda))^{\frac{3}{2}}} = -\frac{s^T(\lambda)\nabla_{\lambda}s(\lambda)}{\|s(\lambda)\|_2^3}. \end{aligned}$$

Differentiating the defining equation

$$(B + \lambda I)s(\lambda) = -g \implies (B + \lambda I)\nabla_{\lambda}s(\lambda) + s(\lambda) = 0.$$

Notice that, rather than $\nabla_{\lambda}s(\lambda)$, merely

$$s^T(\lambda)\nabla_{\lambda}s(\lambda) = -s^T(\lambda)(B + \lambda I)(\lambda)^{-1}s(\lambda)$$

required for $\phi'(\lambda)$. Given the factorization $B + \lambda I = L(\lambda)L^T(\lambda) \implies$

$$\begin{aligned} s^T(\lambda)(B + \lambda I)^{-1}s(\lambda) &= s^T(\lambda)L^{-T}(\lambda)L^{-1}(\lambda)s(\lambda) \\ &= (L^{-1}(\lambda)s(\lambda))^T(L^{-1}(\lambda)s(\lambda)) = \|w(\lambda)\|_2^2 \end{aligned}$$

where $L(\lambda)w(\lambda) = s(\lambda)$.

NEWTON'S METHOD & THE SECULAR EQUATION

Let $\lambda > -\lambda_1$ and $\Delta > 0$ be given

Until "convergence" do:

Factorize $B + \lambda I = LL^T$

Solve $LL^T s = -g$

Solve $Lw = s$

Replace λ by

$$\lambda + \left(\frac{\|s\|_2 - \Delta}{\Delta} \right) \left(\frac{\|s\|_2^2}{\|w\|_2^2} \right)$$

SOLVING THE LARGE-SCALE PROBLEM

- when n is large, factorization may be impossible
- may instead try to use an iterative method to approximate
 - ◊ steepest descent leads to the Cauchy point
 - ◊ obvious generalization: conjugate gradients ... but
 - ▷ what about the trust region?
 - ▷ what about negative curvature?

CONJUGATE GRADIENTS TO “MINIMIZE” $q(s)$

Given $s^0 = 0$, set $g^0 = g$, $d^0 = -g$ and $i = 0$
 Until g^i “small” or breakdown, iterate

$$\alpha^i = \|g^i\|_2^2 / d^{iT} B d^i$$

$$s^{i+1} = s^i + \alpha^i d^i$$

$$g^{i+1} = g^i + \alpha^i B d^i \quad (\equiv g + B s^{i+1})$$

$$\beta^i = \|g^{i+1}\|_2^2 / \|g^i\|_2^2$$

$$d^{i+1} = -g^{i+1} + \beta^i d^i$$

and increase i by 1

CRUCIAL PROPERTY OF CONJUGATE GRADIENTS

Theorem 3.10. Suppose that the conjugate gradient method is applied to minimize $q(s)$ starting from $s^0 = 0$, and that $d^{iT} B d^i > 0$ for $0 \leq i \leq k$. Then the iterates s^j satisfy the inequalities

$$\|s^j\|_2 < \|s^{j+1}\|_2$$

for $0 \leq j \leq k - 1$.

TRUNCATED CONJUGATE GRADIENTS

Apply the conjugate gradient method, but terminate at iteration i if

1. $d^{iT} B d^i \leq 0 \implies$ problem unbounded along d^i
2. $\|s^i + \alpha^i d^i\|_2 > \Delta \implies$ solution on trust-region boundary

In both cases, stop with $s_* = s^i + \alpha^i d^i$, where α^i chosen as positive root of

$$\|s^i + \alpha^i d^i\|_2 = \Delta$$

Crucially

$$q(s_*) \leq q(s^c) \quad \text{and} \quad \|s_*\|_2 \leq \Delta$$

\implies TR algorithm converges to a first-order critical point

HOW GOOD IS TRUNCATED C.G.?

In the convex case ... very good

Theorem 3.11. Suppose that the truncated conjugate gradient method is applied to minimize $q(s)$ and that B is positive definite. Then the computed and actual solutions to the problem, s_* and s_*^M , satisfy the bound

$$q(s_*) \leq \frac{1}{2}q(s_*^M)$$

In the non-convex case ... maybe poor

- e.g., if $g = 0$ and B is indefinite $\implies q(s_*) = 0$
- can use Lanczos method to continue around trust-region boundary if necessary