

Part 2: Linesearch methods for unconstrained optimization

Nick Gould (RAL)

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x)$$

Part C course on continuous optimization

UNCONSTRAINED MINIMIZATION

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x)$$

where the **objective function** $f : \mathbb{R}^n \longrightarrow \mathbb{R}$

- ◉ assume that $f \in C^1$ (sometimes C^2) and Lipschitz
- ◉ often in practice this assumption violated, but not necessary

ITERATIVE METHODS

- in practice very rare to be able to provide explicit minimizer
- iterative method: given starting “guess” x_0 , generate sequence

$$\{x_k\}, \quad k = 1, 2, \dots$$

- **AIM:** ensure that (a subsequence) has some favourable limiting properties:
 - ◊ satisfies first-order necessary conditions
 - ◊ satisfies second-order necessary conditions

Notation: $f_k = f(x_k)$, $g_k = g(x_k)$, $H_k = H(x_k)$.

LINSEARCH METHODS

- calculate a **search direction** p_k from x_k
- ensure that this direction is a **descent direction**, i.e.,

$$g_k^T p_k < 0 \text{ if } g_k \neq 0$$

so that, for small steps along p_k , the objective function **will** be reduced


- calculate a suitable **steplength** $\alpha_k > 0$ so that

$$f(x_k + \alpha_k p_k) < f_k$$

- computation of α_k is the **linsearch**—may itself be an iteration
- generic linsearch method:


$$x_{k+1} = x_k + \alpha_k p_k$$

STEPS MIGHT BE TOO LONG



The objective function $f(x) = x^2$ and the iterates $x_{k+1} = x_k + \alpha_k p_k$ generated by the descent directions $p_k = (-1)^{k+1}$ and steps $\alpha_k = 2 + 3/2^{k+1}$ from $x_0 = 2$

STEPS MIGHT BE TOO SHORT



The objective function $f(x) = x^2$ and the iterates $x_{k+1} = x_k + \alpha_k p_k$ generated by the descent directions $p_k = -1$ and steps $\alpha_k = 1/2^{k+1}$ from $x_0 = 2$

PRACTICAL LINESEARCH METHODS

- in early days, pick α_k to minimize

$$f(x_k + \alpha p_k)$$

- ◊ **exact** linesearch—univariate minimization
 - ◊ rather expensive and certainly not cost effective
 - modern methods: **inexact** linesearch
 - ◊ ensure steps are neither too long nor too short
 - ◊ try to pick “useful” initial stepsize for fast convergence
 - ◊ best methods are either
 - ▷ “backtracking- Armijo” or
 - ▷ “Armijo-Goldstein”
- based

BACKTRACKING LINESEARCH

Procedure to find the stepsize α_k :

Given $\alpha_{\text{init}} > 0$ (e.g., $\alpha_{\text{init}} = 1$)
let $\alpha^{(0)} = \alpha_{\text{init}}$ and $l = 0$
Until $f(x_k + \alpha^{(l)}p_k) \leq f_k$
 set $\alpha^{(l+1)} = \tau\alpha^{(l)}$, where $\tau \in (0, 1)$ (e.g., $\tau = \frac{1}{2}$)
 and increase l by 1
Set $\alpha_k = \alpha^{(l)}$

- this prevents the step from getting too small... but does not prevent too large steps relative to decrease in f
- need to tighten requirement


$$f(x_k + \alpha^{(l)}p_k) \leq f_k$$

ARMJO CONDITION

In order to prevent large steps relative to decrease in f , instead require

$$f(x_k + \alpha_k p_k) \leq f(x_k) + \alpha_k \beta g_k^T p_k$$

for some $\beta \in (0, 1)$ (e.g., $\beta = 0.1$ or even $\beta = 0.0001$)



BACKTRACKING-ARMJO LINESEARCH

Procedure to find the stepsize α_k :

Given $\alpha_{\text{init}} > 0$ (e.g., $\alpha_{\text{init}} = 1$)
let $\alpha^{(0)} = \alpha_{\text{init}}$ and $l = 0$
Until $f(x_k + \alpha^{(l)}p_k) \leq f(x_k) + \alpha^{(l)}\beta g_k^T p_k$
set $\alpha^{(l+1)} = \tau \alpha^{(l)}$, where $\tau \in (0, 1)$ (e.g., $\tau = \frac{1}{2}$)
and increase l by 1
Set $\alpha_k = \alpha^{(l)}$

SATISFYING THE ARMIJO CONDITION

Theorem 2.1. Suppose that $f \in C^1$, that $g(x)$ is Lipschitz continuous with Lipschitz constant $\gamma(x)$, that $\beta \in (0, 1)$ and that p is a descent direction at x . Then the Armijo condition

$$f(x + \alpha p) \leq f(x) + \alpha\beta g(x)^T p$$

is satisfied for all $\alpha \in [0, \alpha_{\max}(x)]$, where

$$\alpha_{\max} = \frac{2(\beta - 1)g(x)^T p}{\gamma(x)\|p\|_2^2}$$

PROOF OF THEOREM 2.1

Taylor's theorem (Theorem 1.1) +

$$\alpha \leq \frac{2(\beta - 1)g(x)^T p}{\gamma(x)\|p\|_2^2},$$

\implies

$$\begin{aligned} f(x + \alpha p) &\leq f(x) + \alpha g(x)^T p + \frac{1}{2}\gamma(x)\alpha^2\|p\|_2^2 \\ &\leq f(x) + \alpha g(x)^T p + \alpha(\beta - 1)g(x)^T p \\ &= f(x) + \alpha\beta g(x)^T p \end{aligned}$$

THE ARMIJO LINESEARCH TERMINATES

Corollary 2.2. Suppose that $f \in C^1$, that $g(x)$ is Lipschitz continuous with Lipschitz constant γ_k at x_k , that $\beta \in (0, 1)$ and that p_k is a descent direction at x_k . Then the stepsize generated by the backtracking-Armijo linesearch terminates with

$$\alpha_k \geq \min \left(\alpha_{\text{init}}, \frac{2\tau(\beta - 1)g_k^T p_k}{\gamma_k \|p_k\|_2^2} \right)$$

PROOF OF COROLLARY 2.2

Theorem 2.1 \implies linesearch will terminate as soon as $\alpha^{(l)} \leq \alpha_{\max}$.
2 cases to consider:

1. May be that α_{init} satisfies the Armijo condition $\implies \alpha_k = \alpha_{\text{init}}$.
2. Otherwise, must be a last linesearch iteration (the l -th) for which

$$\alpha^{(l)} > \alpha_{\max} \implies \alpha_k \geq \alpha^{(l+1)} = \tau \alpha^{(l)} > \tau \alpha_{\max}$$

Combining these 2 cases gives required result.

GENERIC LINESEARCH METHOD

Given an initial guess x_0 , let $k = 0$

Until convergence:

Find a descent direction p_k at x_k

Compute a stepsize α_k using a
backtracking-Armijo linesearch along p_k

Set $x_{k+1} = x_k + \alpha_k p_k$, and increase k by 1

GLOBAL CONVERGENCE THEOREM

Theorem 2.3. Suppose that $f \in C^1$ and that g is Lipschitz continuous on \mathbb{R}^n . Then, for the iterates generated by the Generic Linesearch Method, either

$$g_l = 0 \text{ for some } l \geq 0$$

or

$$\lim_{k \rightarrow \infty} f_k = -\infty$$

or

$$\lim_{k \rightarrow \infty} \min (|p_k^T g_k|, |p_k^T g_k| / \|p_k\|_2) = 0.$$

PROOF OF THEOREM 2.3

Suppose that $g_k \neq 0$ for all k and that $\lim_{k \rightarrow \infty} f_k > -\infty$. Armijo \implies

$$f_{k+1} - f_k \leq \alpha_k \beta p_k^T g_k$$

for all $k \implies$ summing over first j iterations

$$f_{j+1} - f_0 \leq \sum_{k=0}^j \alpha_k \beta p_k^T g_k.$$

LHS bounded below by assumption \implies RHS bounded below. Sum composed of -ve terms \implies

$$\lim_{k \rightarrow \infty} \alpha_k |p_k^T g_k| = 0$$

Let

$$\mathcal{K}_1 \stackrel{\text{def}}{=} \left\{ k \mid \alpha_{\text{init}} > \frac{2\tau(\beta - 1)g_k^T p_k}{\gamma \|p_k\|_2^2} \right\} \quad \& \quad \mathcal{K}_2 \stackrel{\text{def}}{=} \{1, 2, \dots\} \setminus \mathcal{K}_1$$

where γ is the assumed uniform Lipschitz constant.

For $k \in \mathcal{K}_1$,

$$\alpha_k \geq \frac{2\tau(\beta-1)g_k^T p_k}{\gamma \|p_k\|_2^2}$$

\implies

$$\alpha_k p_k^T g_k \leq \frac{2\tau(\beta-1)}{\gamma} \left(\frac{g_k^T p_k}{\|p_k\|} \right)^2 < 0$$

\implies

$$\lim_{k \in \mathcal{K}_1 \rightarrow \infty} \frac{|p_k^T g_k|}{\|p_k\|_2} = 0. \quad (1)$$

For $k \in \mathcal{K}_2$,

$$\alpha_k \geq \alpha_{\text{init}}$$

\implies

$$\lim_{k \in \mathcal{K}_2 \rightarrow \infty} |p_k^T g_k| = 0. \quad (2)$$

Combining (1) and (2) gives the required result.

METHOD OF STEEPEST DESCENT

The search direction

$$p_k = -g_k$$

gives the so-called **steepest-descent** direction.

- ◉ p_k is a descent direction
- ◉ p_k solves the problem

$$\underset{p \in \mathbb{R}^n}{\text{minimize}} \quad m_k^L(x_k + p) \stackrel{\text{def}}{=} f_k + g_k^T p \quad \text{subject to} \quad \|p\|_2 = \|g_k\|_2$$

Any method that uses the steepest-descent direction is a

method of steepest descent.

GLOBAL CONVERGENCE FOR STEEPEST DESCENT

Theorem 2.4. Suppose that $f \in C^1$ and that g is Lipschitz continuous on \mathbb{R}^n . Then, for the iterates generated by the Generic Linesearch Method using the steepest-descent direction,

either

$$g_l = 0 \text{ for some } l \geq 0$$

or

$$\lim_{k \rightarrow \infty} f_k = -\infty$$

or

$$\lim_{k \rightarrow \infty} g_k = 0.$$

PROOF OF THEOREM 2.4

Follows immediately from Theorem 2.3, since

$$\min (|p_k^T g_k|, |p_k^T g_k| / \|p_k\|_2) = \|g_k\|_2 \min (1, \|g_k\|_2)$$

and thus


$$\lim_{k \rightarrow \infty} \min (|p_k^T g_k|, |p_k^T g_k| / \|p_k\|_2) = 0$$

implies that $\lim_{k \rightarrow \infty} g_k = 0$.

METHOD OF STEEPEST DESCENT (cont.)

- archetypical globally convergent method
- many other methods resort to steepest descent in bad cases
- not scale invariant
- convergence is usually very (very!) slow (linear)
- numerically often not convergent at all

STEEPEST DESCENT EXAMPLE



Contours for the objective function $f(x, y) = 10(y - x^2)^2 + (x - 1)^2$, and the iterates generated by the Generic Linesearch steepest-descent method

MORE GENERAL DESCENT METHODS

Let B_k be a symmetric, positive definite matrix, and define the search direction p_k so that

$$B_k p_k = -g_k$$

Then

- p_k is a descent direction
- p_k solves the problem

$$\underset{p \in \mathbb{R}^n}{\text{minimize}} \quad m_k^Q(x_k + p) \stackrel{\text{def}}{=} f_k + g_k^T p + \frac{1}{2} p^T B_k p$$

- if the Hessian H_k is positive definite, and $B_k = H_k$, this is **Newton's method**

MORE GENERAL GLOBAL CONVERGENCE

Theorem 2.5. Suppose that $f \in C^1$ and that g is Lipschitz continuous on \mathbb{R}^n . Then, for the iterates generated by the Generic Linesearch Method using the more general descent direction, either

$$g_l = 0 \text{ for some } l \geq 0$$

or

$$\lim_{k \rightarrow \infty} f_k = -\infty$$

or

$$\lim_{k \rightarrow \infty} g_k = 0$$

provided that the eigenvalues of B_k are uniformly bounded and bounded away from zero.

PROOF OF THEOREM 2.5

Let $\lambda_{\min}(B_k)$ and $\lambda_{\max}(B_k)$ be the smallest and largest eigenvalues of B_k . By assumption, there are bounds $\lambda_{\min} > 0$ and λ_{\max} such that

$$\lambda_{\min} \leq \lambda_{\min}(B_k) \leq \frac{s^T B_k s}{\|s\|^2} \leq \lambda_{\max}(B_k) \leq \lambda_{\max}$$

and thus that

$$\lambda_{\max}^{-1} \leq \lambda_{\max}^{-1}(B_k) = \lambda_{\min}(B_k^{-1}) \leq \frac{s^T B_k^{-1} s}{\|s\|^2} \leq \lambda_{\max}(B_k^{-1}) = \lambda_{\min}^{-1}(B_k) \leq \lambda_{\min}^{-1}$$

for any nonzero vector s . Thus

$$|p_k^T q_k| = |g_k^T B_k^{-1} q_k| \geq \lambda_{\min}(B_k^{-1}) \|q_k\|_2^2 \geq \lambda_{\max}^{-1} \|q_k\|_2^2$$

In addition

$$\|p_k\|_2^2 = g_k^T B_k^{-2} q_k \leq \lambda_{\max}(B_k^{-2}) \|q_k\|_2^2 \leq \lambda_{\min}^{-2} \|q_k\|_2^2,$$

\implies

$$\|p_k\|_2 \leq \lambda_{\min}^{-1} \|q_k\|_2$$

$$\implies \frac{|p_k^T g_k|}{\|p_k\|_2} \geq \frac{\lambda_{\min}}{\lambda_{\max}} \|g_k\|_2$$

Thus

$$\min (|p_k^T g_k|, |p_k^T g_k| / \|p_k\|_2) \geq \frac{\|g_k\|_2}{\lambda_{\max}} \min (\lambda_{\min}, \|g_k\|_2)$$

$$\implies \lim_{k \rightarrow \infty} \min (|p_k^T g_k|, |p_k^T g_k| / \|p_k\|_2) = 0$$

$$\implies \lim_{k \rightarrow \infty} g_k = 0.$$

MORE GENERAL DESCENT METHODS (cont.)

- may be viewed as “scaled” steepest descent
- convergence is often faster than steepest descent
- can be made scale invariant for suitable B_k

CONVERGENCE OF NEWTON'S METHOD

Theorem 2.6. Suppose that $f \in C^2$ and that H is Lipschitz continuous on \mathbb{R}^n . Then suppose that the iterates generated by the Generic Linesearch Method with $\alpha_{\text{init}} = 1$ and $\beta < \frac{1}{2}$, in which the search direction is chosen to be the Newton direction $p_k = -H_k^{-1}g_k$ whenever possible, has a limit point x_* for which $H(x_*)$ is positive definite. Then

- (i) $\alpha_k = 1$ for all sufficiently large k ,
- (ii) the entire sequence $\{x_k\}$ converges to x_* , and
- (iii) the rate is Q-quadratic, i.e, there is a constant $\kappa \geq 0$.

$$\lim_{k \rightarrow \infty} \frac{\|x_{k+1} - x_*\|_2}{\|x_k - x_*\|_2^2} \leq \kappa.$$

PROOF OF THEOREM 2.6

Consider $\lim_{k \in \mathcal{K}} x_k = x_*$. Continuity $\implies H_k$ positive definite for all $k \in \mathcal{K}$ sufficiently large $\implies \exists k_0 \geq 0$:

$$p_k^T H_k p_k \geq \frac{1}{2} \lambda_{\min}(H_*) \|p_k\|_2^2$$

$\forall k_0 \leq k \in \mathcal{K}$, where $\lambda_{\min}(H_*)$ = smallest eigenvalue of $H(x_*) \implies$

$$|p_k^T g_k| = -p_k^T g_k = p_k^T H_k p_k \geq \frac{1}{2} \lambda_{\min}(H_*) \|p_k\|_2^2. \quad (3)$$

$\forall k_0 \leq k \in \mathcal{K}$, and

$$\lim_{k \in \mathcal{K} \rightarrow \infty} p_k = 0$$

since Theorem 2.5 \implies at least one of the LHS of (3) and

$$\frac{|p_k^T g_k|}{\|p_k\|_2} = -\frac{p_k^T g_k}{\|p_k\|_2} \geq \frac{1}{2} \lambda_{\min}(H_*) \|p_k\|_2$$

converges to zero for such k .

Taylor's theorem $\implies \exists z_k$ between x_k and $x_k + p_k$ such that

$$f(x_k + p_k) = f_k + p_k^T g_k + \frac{1}{2} p_k^T H(z_k) p_k.$$

Lipschitz continuity of H & $H_k p_k + g_k = 0 \implies$

$$\begin{aligned} f(x_k + p_k) - f_k - \frac{1}{2} p_k^T g_k &= \frac{1}{2} (p_k^T g_k + p_k^T H(z_k) p_k) \\ &= \frac{1}{2} (p_k^T g_k + p_k^T H_k p_k) + \frac{1}{2} (p_k^T (H(z_k) - H_k) p_k) \\ &\leq \frac{1}{2} \gamma \|z_k - x_k\|_2 \|p_k\|_2^2 \leq \frac{1}{2} \gamma \|p_k\|_2^3 \end{aligned} \tag{4}$$

Now pick k sufficiently large so that

$$\gamma \|p_k\|_2 \leq \lambda_{\min}(H_*) (1 - 2\beta).$$

+ (3) + (4) \implies

$$\begin{aligned} f(x_k + p_k) - f_k &\leq \frac{1}{2} p_k^T g_k + \frac{1}{2} \lambda_{\min}(H_*) (1 - 2\beta) \|p_k\|_2^2 \\ &\leq \frac{1}{2} (1 - (1 - 2\beta)) p_k^T g_k = \beta p_k^T g_k \end{aligned}$$

\implies unit stepsize satisfies the Armijo condition for all sufficiently large

$k \in \mathcal{K}$

Now note that $\|H_k^{-1}\|_2 \leq 2/\lambda_{\min}(H_*)$ for all sufficiently large $k \in \mathcal{K}$. The iteration gives

$$\begin{aligned} x_{k+1} - x_* &= x_k - x_* - H_k^{-1} g_k = x_k - x_* - H_k^{-1} (g_k - g(x_*)) \\ &= H_k^{-1} (g(x_*) - g_k - H_k(x_* - x_k)). \end{aligned}$$

But Theorem 1.3 \implies

$$\|g(x_*) - g_k - H_k(x_* - x_k)\|_2 \leq \gamma \|x_* - x_k\|_2^2$$


\implies

$$\|x_{k+1} - x_*\|_2 \leq \gamma \|H_k^{-1}\|_2 \|x_* - x_k\|_2^2$$

which is (iii) when $\kappa = 2\gamma/\lambda_{\min}(H_*)$. for $k \in \mathcal{K}$.

Result (ii) follows since once iterate becomes sufficiently close to x_* , (iii) for $k \in \mathcal{K}$ sufficiently large implies $k + 1 \in \mathcal{K} \implies \mathcal{K} = \mathbb{N}$. Thus (i) and (iii) are true for all k sufficiently large.

NEWTON METHOD EXAMPLE



Contours for the objective function $f(x, y) = 10(y - x^2)^2 + (x - 1)^2$, and the iterates generated by the Generic Linesearch Newton method

MODIFIED NEWTON METHODS

If H_k is indefinite, it is usual to solve instead

$$(H_k + M_k)p_k \equiv B_k p_k = -g_k$$

where

- M_k chosen so that $B_k = H_k + M_k$ is “sufficiently” positive definite
- $M_k = 0$ when H_k is itself “sufficiently” positive definite

Possibilities:

- If H_k has the spectral decomposition $H_k = Q_k D_k Q_k^T$ then

$$B_k \equiv H_k + M_k = Q_k \max(\epsilon, |D_k|) Q_k^T$$

- $M_k = \max(0, \epsilon - \lambda_{\min}(H_k)) I$
- **Modified Cholesky**: $B_k \equiv H_k + M_k = L_k L_k^T$

QUASI-NEWTON METHODS

Various attempts to approximate H_k :

- Finite-difference approximations:

$$(H_k)e_i \approx h^{-1}(g(x_k + he_i) - g_k) = (B_k)e_i$$

for some “small” scalar $h > 0$

- Secant approximations: try to ensure the **secant condition**

$$B_{k+1}s_k = y_k \approx H_{k+1}s_k, \quad \text{where } s_k = x_{k+1} - x_k \quad \text{and} \quad y_k = g_{k+1} - g_k$$

- **Symmetric Rank-1 method** (but may be indefinite or even fail):

$$B_{k+1} = B_k + \frac{(y_k - B_k s_k)(y_k - B_k s_k)^T}{(y_k - B_k s_k)^T s_k}$$

- **BFGS method**: (symmetric and positive definite if $y_k^T s_k > 0$):

$$B_{k+1} = B_k + \frac{y_k y_k^T}{y_k^T s_k} - \frac{B_k s_k s_k^T B_k}{s_k^T B_k s_k}$$

MINIMIZING A CONVEX QUADRATIC MODEL

For convex models (B_k positive definite)

$$p_k = (\text{approximate}) \arg \min_{p \in \mathbb{R}^n} f_k + p^T g_k^T + \frac{1}{2} p^T B_k p$$

Generic convex quadratic problem: (B positive definite)

$$(\text{approximately}) \min_{p \in \mathbb{R}^n} q(p) = p^T g + \frac{1}{2} p^T B p$$

MINIMIZATION OVER A SUBSPACE

Given vectors $\{d^0, \dots, d^{i-1}\}$, let

- $D^i = (d^0 : \dots : d^{i-1})$
- Subspace $\mathcal{D}^i = \{p \mid p = D^i p_d \text{ for some } p_d \in \mathbb{R}^i\}$
- $p^i = \arg \min_{p \in \mathcal{D}^i} q(p)$

Result: $D^{iT} g^i = 0$, where $g^i = Bp^i + g$

Proof: require $p^i = D^i p_d^i$, where $p_d^i = \arg \min_{p_d \in \mathbb{R}^i} q(D^i p_d)$

But $q(D^i p_d) = p_d^T D^{iT} g + \frac{1}{2} p_d^T D^{iT} B D^i p_d \implies$

$$0 = D^{iT} B D^i p_d^i + D^{iT} g = D^{iT} (B D^i p_d^i + g) = D^{iT} (B p^i + g) = D^{iT} g^i$$

Equivalently: $d^j T g^i = 0$ for $j = 0, \dots, i-1$

MINIMIZATION OVER A SUBSPACE (cont.)

◦ $d^j T g^i = 0$ for $j = 0, \dots, i-1$, where $g^i = Bp^i + g$

Result: $p^i = p^{i-1} - d^{i-1T} g^{i-1} D^i (D^{iT} B D^i)^{-1} e_i$

Proof: Clearly $p^{i-1} \in \mathcal{D}^{i-1} \subset \mathcal{D}^i$

\implies require $p^i = p^{i-1} + D^i p_d^i$, where $p_d^i = \arg \min_{p_d \in \mathbb{R}^i} q(p^{i-1} + D^i p_d)$

$$\begin{aligned} & \text{But } q(p^{i-1} + D^i p_d) \\ &= q(p^{i-1}) + p_d^T D^{iT} (g + Bp^{i-1}) + \frac{1}{2} p_d^T D^{iT} B D^i p_d \\ &= q(p^{i-1}) + p_d^T D^{iT} g^{i-1} + \frac{1}{2} p_d^T D^{iT} B D^i p_d \\ &= q(p^{i-1}) + p_d^T (d^{i-1T} g^{i-1}) e_i + \frac{1}{2} p_d^T D^{iT} B D^i p_d \end{aligned}$$

where e_i is i -th unit vector \implies

$$p_d^i = -d^{i-1T} g^{i-1} (D^{iT} B D^i)^{-1} e_i$$

MINIMIZATION OVER A B-CONJUGATE SUBSPACE

Minimizer over \mathcal{D}^i : $p^i = p^{i-1} - d^{i-1T} g^{i-1} D^i (D^{iT} B D^i)^{-1} e_i$

Suppose in addition the members of \mathcal{D}^i are B -conjugate:

- **B-conjugacy**: $d^{iT} B d^j = 0$ ($i \neq j$)

Result: $p^i = p^{i-1} + \alpha^{i-1} d^{i-1}$, where

$$\alpha^{i-1} = -\frac{d^{i-1T} g^{i-1}}{d^{i-1T} B d^{i-1}}$$

Proof: $D^{iT} B D^i =$ diagonal matrix with entries $d^{jT} B d^j$

for $j = 0, \dots, i-1$

$\implies (D^{iT} B D^i)^{-1} =$ diagonal matrix with entries $1/d^{jT} B d^j$

for $j = 0, \dots, i-1$

$\implies (D^{iT} B D^i)^{-1} e_i = (1/d^{i-1T} B d^{i-1}) e_i$

BUILDING A B-CONJUGATE SUBSPACE

- $d^j T g^i = 0$ for $j = 0, \dots, i - 1$

Since this implies g^i is independent of \mathcal{D}^i , let

$$d^i = -g^i + \sum_{j=0}^{i-1} \beta^{ij} d^j$$

Aim: find β^{ij} so that d^i is B -conjugate to \mathcal{D}^i

Result (orthogonal gradients): $g^i T g^j = 0$ for all $i \neq j$

Proof: $\text{span}\{g^i\} = \text{span}\{d^i\}$

$$\implies g^j = \sum_{k=0}^j \gamma^{j,k} d^k \text{ for some } \gamma^{j,k}$$

$$\implies g^i T g^j = \sum_{k=0}^j \gamma^{j,k} g^i T d^k = 0 \text{ when } j < i$$

BUILDING A B-CONJUGATE SUBSPACE (cont.)

- $d^i = -g^i + \sum_{j=0}^{i-1} \beta^{ij} d^j$
- $d^{jT} g^i = 0$ for $j = 0, \dots, i-1$, where $g^i = Bp^i + g$

Result: $g^{iT} d^i = -\|g^i\|_2^2$

Proof: $g^{iT} d^i = -g^{iT} g^i + \sum_{j=0}^{i-1} \beta^{ij} g^{iT} d^j$

Corollary: $\alpha^i = \frac{\|g^i\|_2^2}{d^{iT} B d^i} \neq 0 \iff g^i \neq 0$

Proof: by definition

$$\alpha^i = -\frac{g^{iT} d^i}{d^{iT} B d^i}$$

BUILDING A B-CONJUGATE SUBSPACE (cont.)

- $d^i = -g^i + \sum_{j=0}^{i-1} \beta^j d^j$
- $g^{iT} g^j = 0$ for all $i \neq j$

Result: $g^{iT} B d^j = 0$ if $j < i - 1$ and $g^{iT} B d^{i-1} = \frac{\|g^i\|_2}{\alpha^{i-1}}$

Proof: $p^{j+1} = p^j + \alpha^j d^j$ & $g^{j+1} = B p^{j+1} + g$

$$\implies g^{j+1} = g^j + \alpha^j B d^j$$

$$\implies g^{iT} g^{j+1} = g^{iT} g^j + \alpha^j g^{iT} B d^j$$

$$\implies g^{iT} B d^j = 0 \text{ if } j < i - 1$$

$$\text{while } g^{iT} g^i = g^{iT} g^{i-1} + \alpha^{i-1} g^{iT} B d^{i-1} \text{ if } j = i - 1$$

$$\implies g^{iT} B d^{i-1} = \|g^i\|_2 / \alpha^{i-1}$$

BUILDING A B-CONJUGATE SUBSPACE (cont.)

- $d^i = -g^i + \sum_{k=0}^{i-1} \beta^{ik} d^k$
- $d^k T B g^i = 0$ if $k < i - 1$ and $d^{i-1 T} B g^i = \|g^i\|_2^2 / \alpha^{i-1}$
- $\alpha^{i-1} = \|g^{i-1}\|_2^2 / d^{i-1 T} B d^{i-1}$

Result: $\beta^{ij} = 0$ for $j < i - 1$ and $\beta^{i\ i-1} \equiv \beta^i = \frac{\|g_i\|_2^2}{\|g_{i-1}\|_2^2}$

Proof: B-conjugacy \implies

$$0 = d^j T B d^i = -d^j T B g^i + \sum_{k=0}^{i-1} \beta^{ik} d^j T B d^k = -d^j T B g^i + \beta^{ij} d^j T B d^j$$

$$\implies \beta^{ij} = d^j T B g^i / d^j T B d^j$$

Result immediate for $j < i - 1$. For $j = i - 1$,

$$\beta^{i\ i-1} = \frac{d^{i-1 T} B g^i}{d^{i-1 T} B d^{i-1}} = \frac{\|g^i\|_2^2}{\alpha^{i-1} d^{i-1 T} B d^{i-1}} = \frac{\|g^i\|_2^2}{\|g^{i-1}\|_2^2}$$

CONJUGATE-GRADIENT METHOD

Given $p^0 = 0$, set $g^0 = g$, $d^0 = -g$ and $i = 0$.

Until g^i “small” iterate

$$\alpha^i = -g^i T d^i / d^i T B d^i$$

$$p^{i+1} = p^i + \alpha^i d^i$$

$$g^{i+1} = g^i + \alpha^i B d^i$$

$$\beta^i = \|g^{i+1}\|_2^2 / \|g^i\|_2^2$$

$$d^{i+1} = -g^{i+1} + \beta^i d^i$$

and increase i by 1

Important features

- $d^j T g^{i+1} = 0 = g^j T g^{i+1}$ for all $j = 0, \dots, i$
- $g^T p^i < 0$ for $i = 1, \dots, n \implies$ descent direction for any $p_k = p^i$
- **stop**: $\|g^i\| \leq \min(\|g\|^\omega, \eta) \|g\|$ ($0 < \eta, \omega < 1$) \implies fast convergence

CONJUGATE GRADIENT METHOD GIVES DESCENT

$$g^{i-1T} d^{i-1} = d^{i-1T} (g + Bp^{i-1}) = d^{i-1T} g + \sum_{j=0}^{i-2} \alpha_j d^{i-1T} B d^j = d^{i-1T} g$$

p^i minimizes $q(p)$ in $\mathcal{D}^i \implies$

$$p^i = p^{i-1} - \frac{g^{i-1T} d^{i-1}}{d^{i-1T} B d^{i-1}} d^{i-1} = p^{i-1} - \frac{g^T d^{i-1}}{d^{i-1T} B d^{i-1}} d^{i-1}.$$

\implies

$$g^T p^i = g^T p^{i-1} - \frac{(g^T d^{i-1})^2}{d^{i-1T} B d^{i-1}},$$

$\implies g^T p^i < g^T p^{i-1} \implies$ (induction)

$$g^T p^i < 0$$

since

$$g^T p^1 = -\frac{\|g\|_2^4}{g^T B g} < 0.$$

$\implies p_k = p^i$ is a descent direction

CG METHODS FOR GENERAL QUADRATICS

Suppose $f(x)$ is quadratic and $x = x_0 + p$

Taylor's theorem \implies

$$f(x) = f(x_0 + p) = f(x_0) + p^T g(x_0) + \frac{1}{2} p^T H(x_0) p$$

- ◉ can minimize as function of p using CG
- ◉ if $x_i = x_0 + p_i \implies g^i = g(x_0) + H(x_0)p_i = g(x_i)$
- ◉ $\alpha^i = -\frac{g(x_i)^T d^i}{d_i^T H(x_0) d^i} = \arg \min_{\alpha} f(x_i + \alpha d^i)$

NONLINEAR CONJUGATE-GRADIENT METHODS

method for minimizing quadratic $f(x)$

Given x^0 and $g(x_0)$, set $d^0 = -g(x_0)$ and $i = 0$.

Until $g(x_k)$ “small” iterate

$$\alpha^i = \arg \min_{\alpha} f(x_i + \alpha d^i)$$

$$x_{i+1} = x_i + \alpha^i d^i$$

$$\beta^i = \frac{\|g(x_{i+1})\|_2^2}{\|g(x_i)\|_2^2}$$

$$d^{i+1} = -g(x_{i+1}) + \beta^i d^i$$

and increase i by 1

may also be used for nonlinear $f(x)$ (Fletcher & Reeves)

- replace calculation of α^i by suitable linesearch
- other methods pick different β^i to ensure descent