


Introduction and Preliminaries

Lecture 1, Continuous Optimisation
Oxford University Computing Laboratory, HT 2006
Notes by Dr Raphael Hauser (hauser@comlab.ox.ac.uk)

Example 1: Linear Programming

A network of gas pipelines is given.



- An arrow from node i to node j represents a pipe with transport capacity c_{ij} in the given direction.
- Transporting one unit of gas along the edge (ij) costs d_{ij} .
- The amount of gas produced at node i is p_i ,
- and the amount of gas consumed is q_i .
- We assume that the total amount consumed equals the total amount of gas produced.
- How to choose the quantities x_{ij} of gas shipped along the edges (ij) so as to minimise costs while satisfying demands?

We set $c_{ij} = 0$ (and d_{ij} arbitrary numbers) for all edges (ij) that do not exist. Doing so, we can assume that the network is a complete graph.

The problem we have to solve is the following:

$$\begin{aligned} \min_x \quad & \sum_{i,j=1}^6 d_{ij}x_{ij} \\ \text{s.t.} \quad & \sum_{k=1}^6 x_{ki} + p_i = \sum_{j=1}^6 x_{ij} + q_i, \quad (i = 1, \dots, 6), \quad (1) \\ & 0 \leq x_{ij} \leq c_{ij}, \quad (i, j = 1, \dots, 6). \quad (2) \end{aligned}$$

- This is an example of a *linear programming* problem, as the *objective function* $\sum_{i,j=1}^6 d_{ij}x_{ij}$ and the *constraint functions* (1),(2) are linear functions of the *decision variables* x_{ij} .
- Note that it is not a priori clear that this problem has feasible solutions. One is therefore interested in algorithms that not only find optimal LP solutions when these exist but also detect when a problem instance is infeasible!
- Furthermore, if there is an optimal solution, we are not only interested in the minimum value of the objective function, but also in the values of x_{ij} that achieve this minimum. Such an x is called a *minimiser* of the problem.

This problem can be modelled as

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \sum_{i=1}^n \sum_{j=1}^n \sigma_{ij} x_i x_j \\ \text{s.t.} \quad & \sum_{i=1}^n \mu_i x_i \geq b, \\ & \sum_{i=1}^n x_i = 1, \\ & x_i \geq 0 \quad (i = 1, \dots, n). \end{aligned}$$

The constraint $\sum_{i=1}^n x_i = 1$ expresses the requirement that 100% of the initial capital has to be invested.

Example 2: Quadratic Programming

- An investor considers a fixed time interval and wishes to decide which fraction of the capital he/she wants to invest in each of n different given assets.
- The expected return of asset i is μ_i , assumed known.
- The covariance between assets i and j is σ_{ij} , assumed known.
- The investor aims at a total return of at least b .
- Subject to this constraint, how to minimise the variance of the overall portfolio (notion of risk)?

Example 3: Semidefinite Programming

- In optimal control, variables y_1, \dots, y_m have to be chosen so as to design a system that is driven by the linear ODE

$$\dot{u} = M(y)u,$$

where $M(y) = \sum_{i=1}^m y_i A_i + A_0$ is an affine combination of given symmetric matrices A_i ($i = 0, \dots, m$).

- To stabilise the system, one would like to choose y so as to minimise the largest eigenvalue of $M(y)$.

Note that $\lambda_1(M) \leq \eta$ if and only if $\eta I - M$ has only non-negative eigenvalues.

This is equivalent to $\eta I - M$ being positive semidefinite, denoted by $\eta I - M \succeq 0$.

The problem we need to solve is thus the following,

$$\begin{aligned} & \max_{\eta, y} -\eta \\ \text{s.t.} \quad & \eta I - A_0 - \sum_{i=1}^m y_i A_i \succeq 0. \end{aligned}$$

This problem can be formulated as

$$\begin{aligned} \text{(P)} \quad & \min_{x, y} x^2 + y^2 - 4 \\ \text{s.t.} \quad & x - 0.5 \geq 0, \\ & -x + 3 \geq 0, \\ & xy - 1 = 0. \end{aligned}$$

Example 4: Polynomial Programming

- An engineer designs a system determined by two design variables x and y which are dependent on each other via the relation $xy = 1$.
- The energy consumed by the system is given by $E(x, y) = x^2 + y^2 - 4$.
- The physical properties of materials used impose the constraints $x \in [0.5, 3]$.
- How to design a system that consumes the smallest amount of energy among all admissible systems?

The General Problem:

More generally, a *continuous programming problem* concerns the minimisation (or maximisation) of a continuous objective function f under constraints defined by continuous functions g_i, h_j :

$$\begin{aligned} \text{(P)} \quad & \min_{x \in \mathbb{R}^n} f(x) \\ \text{s.t.} \quad & g_i(x) \geq 0 \quad (i = 1, \dots, p) \\ & h_j(x) = 0 \quad (j = 1, \dots, q). \end{aligned}$$

What are key properties of iterative algorithms?

- Typically we will assume $f, g_i, h_j \in C^2$.
- $g_i(x) \geq 0$ are called *inequality constraints*.
- $h_j(x) = 0$ are called *equality constraints*.
- Constraints of the form $x_i \in \mathbb{Z}$ (integrality constraints) add a whole other layer of difficulty we will not consider in this course (see Section B course Integer Programming).

- Correct termination: does the algorithm converge to a minimiser? (\rightarrow to recognise optima, need to characterise them mathematically, i.e., develop *optimality conditions*).
- Low complexity:
 - i. low total number of iterations (\rightarrow need a notion of *convergence rate*),
 - ii. low number of computer operations *per* iteration (\rightarrow often leads to a trade-off with i.).
- Reliability: how sensitive is the algorithm to small changes in input, how is it affected by round-off?

Some Terminology:

- $x \in \mathbb{R}^n$ is called *feasible solution* for (P) if it satisfies all the constraints, that is, if

$$\begin{aligned} g_i(x) &\geq 0 & (i = 1, \dots, p), \\ h_j(x) &= 0 & (j = 1, \dots, q). \end{aligned}$$

- The set of feasible solutions is denoted by \mathcal{F} , also called the feasible set. Hence, (P) can be formulated as

$$\min\{f(x) : x \in \mathcal{F}\}.$$

- $x \in \mathbb{R}^n$ is *strictly feasible* if

$$\begin{aligned} g_i(x) &> 0 & (i = 1, \dots, p), \\ h_j(x) &= 0 & (j = 1, \dots, q). \end{aligned}$$

- The set of strictly feasible solutions is denoted by \mathcal{F}° . This is the relative interior of \mathcal{F} .


Example 5: Local versus global optimisation

The problem

$$(P) \quad \min_{x \in \mathbb{R}} f(x) = x^3 + 9x^2$$

$$\text{s.t.} \quad -10 \leq x \leq 2$$

has a local minimiser at $x = 0$, and a global minimiser at $x^* = -10$.



- $x \in \mathcal{F}$ is a *local minimiser* of (P) if there exists $\varepsilon > 0$ such that

$$f(x^*) \leq f(x) \quad \forall x \in \mathcal{F} \cap B_\varepsilon(x^*).$$

- $x^* \in \mathcal{F}$ is a *global minimiser* of (P) if

$$f(x^*) \leq f(x) \quad \forall x \in \mathcal{F}.$$

Q-linear convergence:

- A sequence $(x_k)_{\mathbb{N}} \rightarrow x^* \in \mathbb{R}^n$ converges Q-linearly if there exists $\rho \in (0, 1)$ (the *convergence factor*) and $k_0 \in \mathbb{N}$ such that

$$\|x_{k+1} - x^*\| \leq \rho \|x_k - x^*\| \quad \forall k \geq k_0.$$

- Practical significance: x_k approximates x^* to $O(-\log_{10} \|x_k - x^*\|)$ correct digits. Therefore, $O(-\log_{10} \rho)$ additional correct digits appear per iteration:

$$-\log_{10} \|x_{k+1} - x^*\| - (-\log_{10} \|x_k - x^*\|)$$

$$\geq -\log_{10} \frac{\|x_{k+1} - x^*\|}{\|x_k - x^*\|} \simeq -\log \rho.$$

Example 6:

Let $z \in (0, 1)$ be fixed and consider the sequence $(x_k)_{\mathbb{N}}$ of k -th partial geometric series

$$x_k = \sum_{n=0}^k z^n.$$

Then $(x_k)_{\mathbb{N}}$ converges to $x^* = \frac{1}{1-z} \in \mathbb{R}^1$ Q-linearly with $\rho = z$: for all k ,

$$\frac{|x_{k+1} - x^*|}{|x_k - x^*|} = \frac{\sum_{n=k+2}^{\infty} z^n}{\sum_{m=k+1}^{\infty} z^m} = z < 1.$$

Q-superlinear convergence:

- A sequence $(x_k)_{\mathbb{N}} \rightarrow x^* \in \mathbb{R}^n$ converges Q-superlinearly if

$$\lim_{k \rightarrow \infty} \frac{\|x_{k+1} - x^*\|}{\|x_k - x^*\|} = 0.$$

- Faster than linear for all ρ .
- Practical significance: asymptotically, the number of additional correct digits per iteration becomes larger than any fixed number.

Q-convergence of rate $r > 1$:

- A sequence $(x_k)_{\mathbb{N}} \rightarrow x^* \in \mathbb{R}^n$ converges at the Q-rate $r > 1$ if there exists k_0 such that

$$\|x_{k+1} - x^*\| \leq \|x_k - x^*\|^r \quad \forall k \geq k_0.$$

- Practical significance: the number of additional correct digits is approximately multiplied by r in each iteration:

$$-\log_{10} \|x_{k+1} - x^*\| \simeq r \left(-\log_{10} \|x_k - x^*\| \right).$$

Reading Assignment: Read up on convexity on pages 6–8 of Lecture-Note 1, which can be downloaded from the course web page.