

# The Fundamental Theorem of Linear Inequalities

Lecture 8, Continuous Optimisation

Oxford University Computing Laboratory, HT 2006

Notes by Dr Raphael Hauser ([hauser@comlab.ox.ac.uk](mailto:hauser@comlab.ox.ac.uk))

## Constrained Optimisation and the Need for Optimality Conditions:

In the remaining part of this course we will consider the problem of minimising objective functions over constrained domains. The general problem of this kind can be written in the form

$$\begin{aligned} & \min_{x \in \mathbb{R}^n} f(x) \\ \text{s.t.} \quad & g_i(x) = 0 \quad (i \in \mathcal{E}), \\ & g_j(x) \geq 0 \quad (j \in \mathcal{I}), \end{aligned}$$

where  $\mathcal{E}$  and  $\mathcal{I}$  are the finite index sets corresponding to the equality and inequality constraints, and where  $f, g_i \in C^k(\mathbb{R}^n, \mathbb{R})$  for all  $(i \in \mathcal{I} \cup \mathcal{E})$ .

In unconstrained optimisation we found that we can use the optimality conditions derived in Lecture 1 to transform optimisation problems into zero-finding problems for systems of nonlinear equations

$$\nabla f(x) = 0.$$

We will spend the next few lectures to develop a similar approach to constrained optimisation: in this case the optimal solutions can be characterised by systems of nonlinear equations and inequalities.

A natural by-product of this analysis will be the notion of a *Lagrangian dual* of an optimisation problem: every optimisation problem - called the primal - has a sister problem in the space of Lagrange multipliers - called the dual.

In constrained optimisation it is often advantageous to think of the primal and dual in a combined primal-dual framework where each sheds light from a different angle on a certain saddle-point finding problem.

First we will take a closer look at systems of *linear* inequalities and prove a theorem that will be of fundamental importance in everything that follows:

**Theorem 1: Fundamental theorem of linear inequalities.**

Let  $a_1, \dots, a_m, b \in \mathbb{R}^n$  be a set of vectors. Then exactly one of the two following alternatives occurs:

(I)  $\exists y \in \mathbb{R}_+^m$  such that  $b = \sum_i^m y_i a_i$ .



(II)  $\exists d \in \mathbb{R}^n$  such that  $d^\top b < 0$  and  $d^\top a_i \geq 0$  for all  $(i = 1, \dots, m)$ .

Note that Alternative (I) says that  $b$  lies in the convex cone generated by the vectors  $a_i$ :

$$b \in \text{cone}(a_1, \dots, a_m) := \left\{ \sum_{i=1}^n \lambda_i a_i : \lambda_i \geq 0 \ \forall i \right\}.$$

Alternative (II) on the other hand says that the hyperplane  $d^\perp := \{x \in \mathbb{R}^n : d^\top x = 0\}$  strictly separates  $b$  from the convex set  $\text{cone}(a_1, \dots, a_m)$ .

Thus, Theorem 1 is a result about convex separation: either  $b$  is a member of  $\text{cone}(a_1, \dots, a_m)$  or there exists a hyperplane that strictly separates the two objects.



**Lemma 1:** The two alternatives of Theorem 1 are mutually exclusive.

*Proof:* If this is not the case then we find the contradiction

$$0 \leq \sum_{i=1}^m y_i (d^T a_i) = d^T \left( \sum_{i=1}^m y_i a_i \right) = d^T b < 0. \quad \square$$



**Lemma 2:** W.l.o.g. we may assume that  $\text{span}\{a_1, \dots, a_m\} = \mathbb{R}^n$ .

*Proof:*

- If  $\text{span}\{a_1, \dots, a_m\} \neq \mathbb{R}^n$  then either  $b \in \text{span}\{a_1, \dots, a_m\}$  and then we can restrict all arguments of the proof of Theorem 1 to the linear subspace  $\text{span}\{a_1, \dots, a_m\}$  of  $\mathbb{R}^n$ .
- Else, if  $b \notin \text{span}\{a_1, \dots, a_m\}$  then  $b$  cannot be written in the form  $b = \sum_i^m \mu_i a_i$ , so Alternative (I) does not hold.

- It remains to show that Alternative (II) applies in this case. Let  $\pi$  be the orthogonal projection of  $\mathbb{R}^n$  onto  $\text{span}\{a_1, \dots, a_m\}$ , and let  $d = \pi(b) - b$ . Then  $d \perp \text{span}\{a_1, \dots, a_m\}$ , so that

$$\begin{aligned}d^\top b &= d^\top (b - \pi(b)) + d^\top \pi(b) = -\|d\|^2 + 0 < 0, \\d^\top a_i &= 0 \quad \forall i.\end{aligned}$$

Therefore, Alternative (II) holds. □

Because of Lemma 2, we will henceforth assume that

$$\text{span}\{a_1, \dots, a_m\} = \mathbb{R}^n.$$

We will next construct an algorithm that stops when a situation corresponding to either Alternative (I) or (II) is detected.

This will in fact be the simplex algorithm for LP in disguised form.

## Algorithm 1:

S0 Choose  $J^1 \subseteq \{1, \dots, m\}$  such that  $\text{span}\{a_i\}_{J^1} = \mathbb{R}^n$ ,  $|J^1| = n$ .

S1 For  $k = 1, 2, \dots$  repeat

1. decompose  $b = \sum_{i \in J^k} y_i^k a_i$

2. if  $y_i^k \geq 0 \forall i \in J^k$  return  $y^k$  and stop.

3. else begin

let  $j^k := \min\{i \in J^k : y_i^k < 0\}$

let  $\pi^k : \mathbb{R}^n \rightarrow \text{span}\{a_i : i \in J^k \setminus \{j^k\}\}$  orthogonal projection

let  $d^k := \|a_{j^k} - \pi^k(a_{j^k})\|^{-1}(a_{j^k} - \pi^k(a_{j^k}))$

if  $(d^k)^\top a_i \geq 0$  for  $(i = 1, \dots, m)$  return  $d^k$  and stop.

end

4. let  $l^k := \min\{i : (d^k)^\top a_i < 0\}$

5. let  $J^{k+1} := J^k \setminus \{j^k\} \cup \{l^k\}$

end.

Comments:

- If the algorithm returns  $y^k$  in Step 2, then Alternative (I) holds: let  $y_i = 0$  for  $i \notin J^k$  and  $y_i = y_i^k$  for  $i \in J^k$ . Then  $y \in \mathbb{R}_+^m$  and  $b = \sum_i y_i a_i$ .
- If the algorithm enters Step 3, then  $\{i \in J^k : y_i^k < 0\} \neq \emptyset$  because the condition of Step 2 is not satisfied.
- The vector  $d^k$  constructed in Step 3 satisfies

$$(d^k)^\top b = \sum_{i \in J^k} y_i^k (d^k)^\top a_i = y_{j^k}^k (d^k)^\top a_{j^k} < 0, \quad (1)$$

Therefore, if the algorithm returns  $d^k$  then Alternative (II) holds with  $d = d^k$ .

- If the algorithm enters Step 4 then  $\{i : (d^k)^\top a_i < 0\} \neq \emptyset$  because the condition of the last “if” statement of Step 3 is not satisfied. Moreover, since

$$\begin{aligned} (d^k)^\top a_{j^k} &= 1, \\ (d^k)^\top a_i &= 0 \quad (i \in J^k \setminus \{j^k\}), \end{aligned}$$

we have  $\{i : (d^k)^\top a_i < 0\} \cap J^k = \emptyset$ . This shows that  $l^k \notin J^k$ .

- We have  $\text{span}\{a_i : i \in J^{k+1}\} = \mathbb{R}^n$ , because  $(d^k)^\top a_{l^k} \neq 0$  and  $(d^k)^\top a_i = 0$  ( $i \in J^k \setminus \{j^k\}$ ) show that  $a_{l^k} \notin \text{span}\{a_i : i \in J^k \setminus \{j^k\}\}$ . Moreover,  $|J^{k+1}| = n$ .

**Lemma 3:** It can never occur that  $J^k = J^t$  for  $k < t$ .

*Proof:*

- Let us assume to the contrary that  $J^k = J^t$  for some iterations  $k < t$ , and let  $j^{\max} := \max\{j^s : k \leq s \leq t - 1\}$ .
- Then there exists  $p \in \{k, k + 1, \dots, t - 1\}$  such that  $j^{\max} = j^p$ .
- Since  $J^k = J^t$ , there also exists  $q \in \{k, k + 1, \dots, t - 1\}$  such that  $j^{\max} = j^q$ . In other words, the index must have once left  $J$  and then reentered, or else it must have entered and then left again.



- Now  $j^{\max} = j^p$  implies that for all  $i \in J^p$  such that  $i < j^{\max}$  we have  $y_i^p \geq 0$ .

- Likewise, for the same indices  $i$  we have  $(d^q)^\top a_i \geq 0$ , as

$$i < j^{\max} = l^q = \min\{i : (d^q)^\top a_i < 0\}.$$

- Furthermore, we have  $y_{j^{\max}}^p = y_{j^p}^p < 0$  and  $(d^q)^\top a_{j^{\max}} = (d^q)^\top a_{l^q} < 0$ .

- And finally, since  $J^s \cap \{j^{\max} + 1, \dots, m\}$  remains unchanged for  $s = k, \dots, t$  we have  $(d^q)^\top a_i = 0$  for all  $i \in J^p$  such that  $i > j^{\max}$ .

- Therefore,

$$(d^q)^\top b = \sum_{i \in J^p} y_i^p (d^q)^\top a_i \geq 0. \quad (2)$$

- On the other hand, (1) shows  $(d^q)^\top b < 0$ , contradicting (2). Thus, indices  $k < t$  such that  $J^k = J^t$  do not exist.  $\square$

We are finally ready to prove the fundamental theorem of linear inequalities:

*Proof: (Theorem 1)*

- Since  $J^k \subseteq \{1, \dots, m\}$  and there are finitely many choices for these index sets and Lemma 3 shows that there are no repetitions in the sequence  $J^1, J^2, \dots$ , the sequence must be finite.
- But this is only possible if in some iteration  $k$  Algorithm 1 either returns  $y^k$ , detecting that Alternative (I) holds, or  $d^k$ , detecting that Alternative (II) holds.  $\square$

## The Implicit Function Theorem:

Another fundamental tool we will need is the implicit function theorem.

Before stating the theorem, let us illustrate it with an example:

**Example 1:** The function  $f(x_1, x_2) = x_1^2 + x_2^2 - 1$  has a zero at the point  $(1, 0)$  and  $\frac{\partial}{\partial x_1} f(1, 0) = 1 \neq 0$ . In a neighbourhood of this point the level set  $\{(x_1, x_2) : f(x_1, x_2) = 0\}$  can be explicitly parameterised in terms of  $x_2$ , that is, there exists a function  $h(t)$  such that  $f(x_1, x_2) = 0$  if and only if  $(x_1, x_2) = (h(t), t)$  for some value of  $t$ .

- The level set is nothing else but the unit circle  $S^1$ , and for  $(x_1, x_2)$  with  $x_1 > 0$  we have  $f(x_1, x_2) = 0$  if and only if  $x_1 = h(x_2)$  where  $h(t) = \sqrt{1 - t^2}$ .

- Thus,

$$S^1 \cap \{x \in \mathbb{R}^2 : x_1 > 0\} = \{(h(t), t) : t \in (-1, 1)\},$$





as claimed.

- Another way to say this is that  $S^1$  is a differentiable manifold with local coordinate map

$$\begin{aligned} \varphi : S^1 \cap \{x \in \mathbb{R}^2 : x_1 > 0\} &\rightarrow (-1, 1), \\ x &\mapsto x_2. \end{aligned}$$

- The parameterisation in terms of  $x_2$  was only possible because  $\frac{\partial}{\partial x_1} f(1, 0) \neq 0$ .
- To illustrate this, note that we also have  $f(0, 1) = 0$ , but now  $\frac{\partial}{\partial x_1} f(0, 1) = 0$  and we cannot parameterise  $S^1$  by  $x_2$  in a neighbourhood of  $(0, 1)$ .
- In fact, in a neighbourhood of  $x_2 = 1$ , there are two  $x_1$  values  $\pm\sqrt{1 - x_2^2}$  such that  $f(x_1, x_2) = 0$  when  $x_2 < 1$  and none when  $x_2 > 0$ .

## Example 2:



*Generalisation:*

- $f \in C^k(\mathbb{R}^{p+q}, \mathbb{R}^p)$ .
- Let  $f'_B(x)$  be the leading  $p \times p$  block of the Jacobian matrix  $f'(x) = [f'_B(x) \ f'_N(x)]$ ,
- and  $f'_N(x)$  the trailing  $p \times q$  block.
- Let  $x_B$  be the first  $p \times 1$  block of the vector  $x$
- and  $x_N$  the trailing  $q \times 1$  block.



**Theorem 2: Implicit Function Theorem.** Let  $f \in C^k(\mathbb{R}^{p+q}, \mathbb{R}^p)$  and let  $\bar{x} \in \mathbb{R}^{p+q}$  be such that  $f(\bar{x}) = 0$  and  $f'_B(\bar{x})$  nonsingular.

Then there exist open neighbourhoods  $U_B \subset \mathbb{R}^p$  of  $\bar{x}_B$  and  $U_N \subset \mathbb{R}^q$  of  $\bar{x}_N$  and a function  $h \in C^k(U_N, U_B)$  such that for all  $(x_B, x_N) \in U_B \times U_N$ ,

i)  $f(x_B, x_N) = 0 \Leftrightarrow x_B = h(x_N)$ ,

ii)  $f'_B(x)$  is nonsingular,

iii)  $h'(x_N) = -\left(f'_B(x)\right)^{-1} f'_N(x)$ .

**Reading Assignment:** Lecture-Note 8.